

TYLOSIN 200

COMPOSITION:

Tylosin tartrate eq. 200 mg base
Excipients up to 1 ml.

PHARMACOLOGICAL PROPERTIES:

Macrolide antibiotic active against Gram-positive bacteria, some Gram-negative bacteria, spirochetes (e.g. *Treponema hyodysenteriae*), and mycoplasma.

For the treatment of mycoplasmosis Tylosin 200 injectable solution is the drug of choice.

INDICATIONS:

Horses, camels, cattle, sheep, goats, pigs, dogs, cats, poultry.

Prevention and treatment of:

- respiratory infections in all animals such as broncho-pneumonia, rhinitis, enzootic pneumonia and atrophic rhinitis in pigs.
- mycoplasma infections in poultry (CRD, infectious sinusitis), pigs (enzootic pneumonia, arthritis), and in sheep, goats and camel.
- wound- and skin infections – metritis, mastitis, foul-in-the-foot in cattle – erysipelas, dysentery in pigs – leptospirosis – tonsillitis, otitis externa in dogs – secondary infections.

DOSAGE AND ADMINISTRATION:

By deep I.M. injection or subcutaneous injection (dogs, cats, poultry).

Dose:

10 - 15 mg tylosin per kg body weight or 5 - 7.5 ml Tylosin 200 per 100 kg body weight.

Poultry: 25 mg per kg b.w., once a day, subcutaneously.

If necessary, injections should be repeated with 12 hours intervals.

Duration of treatment:

2 to 5 days for dogs and cats.

1 to 3 days for ruminants and horses.

Once in poultry.

WARNINGS:

Irritations at the injection site and pain occasionally may be observed.

Allergic reactions only seldom occur.

PACK SIZE:

100 ml.

