

Drug Chart

$$\text{Volume (mL)} = (\text{dose (mg/kg)} \times \text{weight (kg)}) \div \text{concentration (mg/mL)}$$

Drug	Concentration (mg/ml)	Dose (mg/kg)	Example Calculation (Goat weighing 50 kg)	Route of Administration	Indication	Contraindications	Withdrawal Time
Flunixin meglumine (NSAID)	50mg/ml	2.2mg/kg	$(2.2\text{mg/kg} \times 50\text{kg}) / 50\text{mg/ml} = 2.2\text{ml}$	IV, IM, Orally	<ul style="list-style-type: none"> Treatment of pain, inflammation and swelling. 	<ul style="list-style-type: none"> Use with caution in animals with cardiac or hepatic impairment. Do not use if parturition is expected within 48 hours 	Meat - 5-14 d Milk - 12-48 hrs
Meloxicam (NSAID)	120mg/ml	2mg/kg	$2\text{mg/kg} \times 50\text{kg} / 120\text{mg/ml} = 0.8\text{ml}$	Orally	<ul style="list-style-type: none"> Given SC or IV for pain relief and inflammation 	<ul style="list-style-type: none"> Ulcerated GIT Nephrotoxicity Not to be used in pregnant animals 	Meat - 15 days Milk - 5 days

Procaine Penicillin G (Antibiotic)	200,000IU	20,000IU	$(2\text{mg/kg} \times 50\text{kg}) / 20 = 5\text{ml}$	IM	<ul style="list-style-type: none"> Prevents infection from Gram negative bacteria such as <i>Streptococci</i>, <i>Pasteurella multocida</i> and <i>Mannheimia haemolytica</i> 	<ul style="list-style-type: none"> Avoid injection volumes greater than 30 mL per site. 	30 d
Lidocaine hydrochloride (Local anesthetic)	20mg/ml (2%)	2mg/kg	$(2\text{mg/kg} \times 50\text{kg}) / 20\text{mg/ml} = 5\text{ml}$	IV, SC, Epidural, Topical	<ul style="list-style-type: none"> 100% nerve block. Short acting local anesthetic. Effective for 45 to 90 minutes 	<ul style="list-style-type: none"> Use with caution in animals that are pregnant and those with cardiac or hepatic impairment 	Meat - 1 day Milk - 24 hours
Xylazine (Alpha 2 agonist)	20mg/ml (2%)	0.05-0.2mg/kg	$(0.05\text{mg/kg} \times 50\text{kg}) / 20\text{mg/ml} = 0.1\text{ml}$	IV, IM	<ul style="list-style-type: none"> Preanesthetic Analgesia for a few hours 	<ul style="list-style-type: none"> Cardiac or respiratory disease Hypotensive animals Maintain in sternal 	Meat - 14 d Milk - 48hr

						recumbency to avoid bloat	
Tolazoline	100mg/ml	0.1mg/kg	(0.1mg/kg x 50kg) / 100mg/ml = 0.05ml	IV, IM	<ul style="list-style-type: none"> • Reversal agent for Xylazine 		4 days
Tetanus antitoxin	15000 IU/ml	5000 IU/ml		IV			4 days

References

- 1) Ezproxy Login [Internet]. Www-vetstream-com.ezproxy.sastudents.uwi.tt. 2020 [cited 2 November 2020]. Available from: <https://www-vetstream-com.ezproxy.sastudents.uwi.tt/treat/bovis/generics/lidocaine>
- 2) Penicillin Injectable for Animal Use [Internet]. Drugs.com. [cited 2020Nov2]. Available from: <https://www.drugs.com/vet/penicillin-injectable.html>
- 3) Drugs for goats [Internet]. Wiley Online Library. John Wiley & Sons, Ltd; 2016 [cited 2020Nov16]. Available from: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/9781119073543.app1>
- 4) 2. Anesthesia Guidelines: Ruminants [Internet]. AHC Research Services - University of Minnesota. 2020 [cited 16 November 2020]. Available from: <https://www.researchservices.umn.edu/services-name/research-animal-resources/research-support/guidelines/anesthesia-ruminants>